


FRANZ^A

Winterard A-quality for light soils



Advantages:

- Very high yielding A-variety with good winterhardiness and stable hagberg
- Very stress tolerant
- For early sowing dates and as stubble wheat

Cultivation:

Adequate and in time doses of plant growth regulators -(splitting)

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development



Vitality

Winter hardiness (6) medium to high

Drought resistance (7) high



Plant Health



Quality



FRANZ^A

Winterard A-quality for light soils

Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	(6) medium to high
Drought resistance	(7) high

Yield structure:

Yield structure	Ear density type
Ears/m ²	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Sowing:

Time of sowing	early until optimum
Seed rate	locally adapted

Straw:

Plant height	(5) medium
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	significant more

Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
DTR	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Eye spot	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Fungicides for intensive cultivation:

Early treatment	according standard
Final treatment	determined application of fungicides (BBCH 59 - 65)

FRANZ ^A

Winterard A-quality for light soils

Quality:

Hagberg / Falling number	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hagberg stability	strong
Zeleny / Sedimentation value	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Protein content	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Water absorption	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Flour extraction	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Loaf volume	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>