


TIFFANY

Low level of vicin and convicin are especially suitable for laying hens and poultry feeding



Advantages:

- Above-average protein yields with very low Vicin and Convicin Contents
- This special quality is interesting for poultry feeding
- High yielding field bean with medium to early maturity and good standing power

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Lodging resistance	
Beginning of flowering	
Yield potential	
Protein content	

TIFFANY

Low level of vicin and convicin are especially suitable for laying hens and poultry feeding

Growth:

Grain maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant length	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Beginning of flowering	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Resistance:

Lodging resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Botrytis	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Yield & Quality:

Yield potential	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein yield	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Thousand grain weight	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Tannin content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Cultivation:

Sowing depth	Heavy soil: 6-8 cm, light soil: 8-10 cm
Plant density	35-50 plants per m ² . The best possible row distance is 25-45 cm depending on the sowing method.
Sowing time	Depending on climatic and soil conditions, beginning at the end of February.