

# FAUSTUS<sup>B</sup>

High-yielding, early maturity, sowing flexibility



## Advantages:

- Beautiful high-yielding variety with earlier heading and maturity
- Suitable before rape and challenging catch crops
- High standing power, very healthy, high and stable falling number

## Cultivation:

Suitable for all previous crop and all regions with fast ripening conditions. Late harvest possible

## Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

### Plant Development



### Vitality

Winter hardiness	(5) medium
Drought resistance	(6) medium to high
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

### Plant Health



### Quality



# FAUSTUS<sup>B</sup>

High-yielding, early maturity, sowing flexibility

## Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	(5) medium
Drought resistance	(6) medium to high

## Yield structure:

Yield structure	ear density type
Ears/m <sup>2</sup>	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

## Sowing:

Time of sowing	all sowing dates
Seed rate	locally adapted

## Straw:

Plant height	medium
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	lower

## Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
DTR	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Eye spot	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

# FAUSTUS<sup>B</sup>

High-yielding, early maturity, sowing flexibility

## Quality:

---

Hagberg / Falling number	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hagberg stability	strong
Zeleny / Sedimentation value	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Protein content	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Water absorption	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Flour extraction	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Loaf volume	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>