

ANAPOLIS^c



Advantages:

- High yielding feed wheat with good lodging resistance
- Excellent disease resistance (mildew, yellow rust)
- Low fusarium risk

Cultivation:

Prefer growing of ANAPOLIS in winter mild regions, suitable in rotations with ALL previous crops, even maize

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development



Vitality

Winter hardiness medium to low

Drought resistance medium



Plant Health



Quality



ANAPOLIS^c

Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	medium to low
Drought resistance	medium

Yield structure:

Yield structure	compensation type
Ears/m ²	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Sowing:

Time of sowing	optimum until late
Seed rate	locally adapted

Straw:

Plant height	short
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	significant lower

Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
DTR	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Fungicides for intensive cultivation:

Early treatment	according standard
Final treatment	according standard

ANAPOLIS^c

Quality:

Hagberg / Falling number	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hagberg stability	weak
Zeleny / Sedimentation value	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Protein content	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Water absorption	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Flour extraction	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Loaf volume	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>