

HIVERNA

Winter field bean with a good winterhardiness



Advantages:

- Winter field bean with a good winterhardiness
- Very early flower und medium maturity
- Significant yield advantage possible

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Lodging resistance	■	■	■	■	■	■	■	■	■
Beginning of flowering	■	■	■	■	■	■	■	■	■
Yield potential	■	■	■	■	■	■	■	■	■
Protein content	■	■	■	■	■	■	■	■	■

HIVERNA

Winter field bean with a good winterhardiness

Growth:

Grain maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant length	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Beginning of flowering	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Resistance:

Lodging resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Ascochyta	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Botrytis	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Yield & Quality:

Yield potential	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Thousand grain weight	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Tannin content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Cultivation:

Sowing depth	Heavy soil: 6-8 cm, light soil 8-10 cm
Plant density	The optimum plant density must not exceed 20 to max. 25 plants per m2 before winter. The optimal distance between rows is 25-45 cm depending on sowing method.
Sowing time	Depending on climatic and soil conditions, between 15th and 30th of September.