

TAIFUN

Good agronomical traits and high protein digestibility for feeding of monogastric



Advantages:

- Field bean with early flowering date
- Free of tannins
- Interesting for the feeding of monogastric animals

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Lodging resistance	■	■	■	■	■	■	■	■	■
Beginning of flowering	■	■	■	■	■	■	■	■	■
Yield potential	■	■	■	■	■	■	■	■	■
Protein content	■	■	■	■	■	■	■	■	■

TAIFUN

Good agronomical traits and high protein digestibility for feeding of monogastric

Growth:

Grain maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant length	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Beginning of flowering	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Resistance:

Lodging resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Botrytis	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Yield & Quality:

Yield potential	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein yield	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Thousand grain weight	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Tannin content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Cultivation:

Sowing depth	Heavy soil: 6-8 cm, light soil: 8-10 cm
Plant density	35-50 plants per m ² . The best possible row distance is 25-45 cm depending on the sowing method.
Sowing time	Depending on climatic and soil conditions, beginning at the end of February.