

QUINTUS^A

Awned, medium early, good disease resistance



Advantages:

- Top 1 yield level in almost all official and postregistration trials in Europe
- High resistance against fusarium, yellow rust and brown rust
- Registration in 5 countries, including Germany, Baltic states and Ukraine

Cultivation:

Cultivation highly profitable due to high yield and low fungicide intensity
Late autumn sowing in winter mild regions possible

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development



Vitality

Drought resistance (7) high



Plant Health



Quality



QUINTUS^A

Awned, medium early, good disease resistance

Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Drought resistance	(7) high

Yield structure:

Yield structure	Single ear type
Ears/m ²	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Sowing:

Time of sowing	middle of February till begin of April
Seed rate	locally adapted

Straw:

Plant height	(4) medium to short
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	slightly lower

Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

QUINTUS ^A

Awned, medium early, good disease resistance

Quality:

Hagberg / Falling number	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Zeleny / Sedimentation value	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Water absorption	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Flour extraction	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Loaf volume	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>