

MATTHUS^A

Alternative wheat



Advantages:

- Alternative wheat with short straw and high winterhardiness
- Clearly improved yields compared to concurrently tested winterwheat
- Best commercial quality (protein, falling number, zeleny) high to very high

Cultivation:

Premium wheat production after late harvest crops. Wide sowing time from middle of October to April.

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development



Vitality

Winter hardiness medium

Drought resistance high



Plant Health



Quality



MATTHUS^A

Alternative wheat

Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	medium
Drought resistance	high

Yield structure:

Yield structure	Ear density type
Ears/m ²	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Sowing:

Time of sowing	15th of October till end of March
Seed rate	locally adapted

Straw:

Plant height	short
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	standard

Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

Fungicides for intensive cultivation:

Early treatment	Treatment against eyespot and septoria after autumn sowing in EC 31/32
Final treatment	combination product in EC 47/51

MATTHUS _A

Alternative wheat

Quality:

Hagberg / Falling number	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Zeleny / Sedimentation value	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Protein content	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Water absorption	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Flour extraction	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Loaf volume	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>