

SU HYMALAYA ^A

Advantages:

- Excellent yield
- Good ear and leaves diseases resistance
- High winter hardiness
- Early sowing for northern countries

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development

Maturity



Vitality

Winter hardiness

8 very good

Drought resistance

high

Lodging resistance



Plant Health

Fusarium



Quality

Hagberg / Falling number



Protein content



SU HYMALAYA^A

Plant development and vitality:

| | |
|--------------------|---------------------------------------------------------------------------------------------------------------------------|
| Ear emergence | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Maturity | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Winter hardiness | 8 very good |
| Drought resistance | high |

Sowing:

| | |
|----------------|-----------------------|
| Time of sowing | very early to optimum |
| Seed rate | 120-140 |

Straw:

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------------------|
| Plant height | medium |
| Lodging resistance | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Plant growth regulator | locally adapted slightly more |

Disease resistance:

| | |
|----------------------|---------------------------------------------------------------------------------------------------------------------------|
| Mildew | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Septoria leaf blotch | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Yellow rust | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Brown rust | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| DTR | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Fusarium | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |

Fungicides for intensive cultivation:

| | |
|-----------------|----------------------------|
| Early treatment | Only occasionally required |
| Final treatment | Standard |

SU HYMALAYA ^A

Quality:

| | |
|------------------------------|----------------------------------------------------------------------------------------------------------------|
| Hagberg / Falling number | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Hagberg stability | Good |
| Zeleny / Sedimentation value | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Protein content | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Water absorption | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Flour extraction | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |
| Loaf volume | <div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div> |